

hep-th/9908198
August 1999

Dilatonic AdS-Kerr Solution to AdS/CFT Correspondence

Alexander BURINSKII¹

NSI Russian Academy of Science,
B. Tulkaya 52, Moscow 113191, RUSSIA

Abstract

We consider the AdS_5 solution deformed by a non-constant dilaton interpolating between the standard AdS (UV region) and flat boundary background (IR region). We show that this dilatonic solution can be generalized to the case of a non-flat boundaries provided that the metric of the boundaries satisfies the vacuum Einstein field equations.

As an example, we describe the case when the four-dimensional boundaries represent the Kerr space-time.

¹ e-mail : grg@ibrae.ac.ru

The recently discovered recently remarkable AdS/CFT correspondence between higher-dimensional classical supergravity and quantum gauge theory on the boundary (bulk/boundary correspondence) [1] gives a new insight to the understanding of strongly coupled gauge theories. In particular, the $AdS_5 \times S^5$ vacua of type-IIB supergravity correspond to the four-dimensional Super Yang Mills theory on the boundary. This stimulated attempts to get the description of running gauge coupling of Yang-Mills theory and QCD-confinement in the frames of type-0 superstring theory [2, 3, 4]. In another approach [5, 6, 7, 8, 9], the attempts to reproduce similar QCD-effects were based on the non-supersymmetric background solutions of type-IIB string theory which can be obtained due to the deformation of $AdS_5 \times S^5$ vacuum by a non-constant dilaton breaking the supersymmetry and conformal symmetry. The exact solution of a such sort was first given in [10] and we shall follow the notations of this work. Starting with ten-dimensional dilatonic gravity and the solution with topology of $AdS_5 \times S^5$, one can integrate out five coordinates of the sphere S^5 and obtain [10, 9] effective action for AdS_5 background

$$S = - \int d^5x \sqrt{-g} \left(R - \Lambda - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \right) , \quad (1)$$

and the equations for metric g and dilaton field ϕ

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + \frac{\Lambda}{2} g_{\mu\nu} - \frac{1}{2} \left(\partial_\mu \phi \partial_\nu \phi - \frac{1}{2} g_{\mu\nu} g^{\rho\sigma} \partial_\rho \phi \partial_\sigma \phi \right) = 0, \quad (2)$$

$$\partial_\mu \left(\sqrt{-g} g^{\mu\nu} \partial_\nu \phi \right) = 0 . \quad (3)$$

Assuming that solutions for g and ϕ depend only on coordinate $y \equiv x^5$, the following ansatz for metric was proposed

$$ds^2 = \sum_{\mu, \nu=0}^d g_{\mu\nu} dx^\mu dx^\nu = f(y) dy^2 + g(y) \sum_{i,j=0}^{d-1} \eta_{ij} dx^i dx^j, \quad (4)$$

where η_{ij} is the metric of Minkowski background. As it was shown in [10, 9], the solution of these equations can be given in five dimensions by functions

$$g = y \quad (5)$$

and

$$f = \frac{3}{y^2 \left(\lambda^2 + \frac{c^2}{2y^4} \right)} , \quad (6)$$

where $\lambda^2 = -\Lambda$ is positive. The main peculiarity of this solution is the appearance of two boundaries corresponding to different limiting values of the dilaton. This solution interpolates between conformal AdS background (weak coupling regime, $y \rightarrow \infty$) and flat space at singular value of dilaton $\phi \rightarrow 6^{\frac{1}{2}} \ln y$ (strong coupling regime, $y \rightarrow 0$).

The aim of this note is to attract attention to the possibility of the generalization of this solution for non-flat four-dimensional boundaries. We shall show that any metric of the form

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = f(y) dy^2 + g(y) \hat{g}_{ik} dx^i dx^k \quad (7)$$

represents the solution of dilatonic gravity equations (2) and (3) provided that the boundary metric $\hat{g}_{ik}$ satisfies the vacuum Einstein equations. This result is also valid for a d-dimensional boundary of AdS_{d+1} . As an example we consider in more details d=4 case when the four-dimensional boundary represents the Kerr geometry.

Let us consider d+1-dimensional metric (7) where $\hat{g}_{ik}$ is a d-dimensional boundary metric, $\mu, \nu \dots = 0, 1, \dots, d$, and $i, k, \dots = 0, 1, \dots, d-1$. In the matrix notations we have

$$g_{\mu\nu} = \begin{pmatrix} g(y) \hat{g}_{ik} & 0 \\ 0 & f \end{pmatrix}, \quad (8)$$

while the contravariant form of this metric is given by

$$g^{\mu\nu} = \begin{pmatrix} g(y)^{-1} \hat{g}^{ik} & 0 \\ 0 & f^{-1} \end{pmatrix}, \quad (9)$$

where $\hat{g}^{ik}$ is the contravariant form of the corresponding d-dimensional metric. The expressions for the connection coefficients $\Gamma_{\nu\lambda}^\mu$ are given in Appendix. For the Ricci curvature tensor we have

$$R_{dd} = -d \left(\frac{g''}{2g} + \left(\frac{g'}{2g} \right)^2 + \frac{f'g'}{4fg} \right), \quad (10)$$

$$R_{id} = 0, \quad (11)$$

$$R_{ik} = \hat{R}_{ik} + g_{ik} \left[\frac{f'g'}{4f^2} - \frac{g''}{2f} + (2-d) \frac{(g')^2}{4fg} \right]. \quad (12)$$

The important point is that $\hat{R}_{ik}$ represents Ricci curvature of d-dimensional boundary with the metric $\hat{g}_{ik}$ and the connections $\hat{\Gamma}_{ik}^l$. Similarly, the scalar curvature is

$$R = g^{-1} \hat{R} + \frac{d}{4f} \left[2 \frac{f'g'}{fg} - 4 \frac{g''}{g} + (3-d) \left(\frac{g'}{g} \right)^2 \right]. \quad (13)$$

Finally, the expressions for the Einstein tensor are given by $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R$ and are as follows:

$$G_{dd} = \frac{d(d-1)}{8}\left(\frac{g'}{g}\right)^2, \quad (14)$$

$$G_{id} = 0, \quad (15)$$

$$G_{ik} = \hat{G}_{ik} + g_{ik}\left[\frac{(d-1)(d-4)(g')^2}{8fg} - \frac{(d-1)f'g'}{4f^2} + \frac{(d-1)g''}{2f}\right]. \quad (16)$$

When the d-dimensional metric $\hat{g}_{ik}$ satisfies the vacuum Einstein field equations $\hat{G}_{ik} = 0$, the components $\hat{G}_{ik}$ drop out of the equations (2) and (3). As a result we come exactly to differential equations for functions f and g given in the paper [9] and to the expressions (5) and (6). Therefore, in the solutions (4), (5) and (6) the d-boundary metric η_{ik} can be replaced by any metric $\hat{g}_{ik}$ satisfying the vacuum Einstein equations.

As an example let us now consider the partial case $d = 4$ and the Kerr boundary metric $\hat{g}_{ik}$ in the Kerr-Schild form

$$\hat{g}_{ik} = \eta_{ik} + 2hk_i k_k, \quad (17)$$

where h is the harmonic scalar function

$$h = mr/(r^2 + a^2 \cos^2 \theta), \quad (18)$$

and m is the mass parameter. This is an important solution of the vacuum Einstein equations describing the field of rotating black holes and modelling the gravitational field of spinning particle [11, 12]. Besides, this form allows one to get a simple comparison with the Minkowskian case.

The vector field $k_i(x)$ is tangent to the principal null congruence and it is determined by the 1-form

$$k_i dx^i = \frac{\sqrt{2}}{1 + Y\bar{Y}}[du + \bar{Y}d\zeta + Yd\bar{\zeta} - Y\bar{Y}dv], \quad (19)$$

where

$$\begin{aligned} 2^{\frac{1}{2}}\zeta &= x + iy, & 2^{\frac{1}{2}}\bar{\zeta} &= x - iy, \\ 2^{\frac{1}{2}}u &= z + t, & 2^{\frac{1}{2}}v &= z - t, \end{aligned} \quad (20)$$

are the Cartesian null coordinates, and $Y(x) = e^{i\phi} \tan \frac{\theta}{2}$ is the projective angular coordinate.² The field k_i is null in respect to Minkowski metric, $k_i k_k \eta^{ik} = 0$, as well as regarding the metric $\hat{g}_{ik}$

$$k_i k_k \hat{g}^{ik} = 0. \quad (21)$$

One can build the five dimensional field $k^\mu = (k^i, 0)$ which will obviously be the null field with respect to the full five-dimensional metric as well

$$k^\mu k^\nu g_{\mu\nu} = 0. \quad (22)$$

Therefore, the five-dimensional Kerr-AdS metric takes the form

$$ds^2 = f(y)dy^2 + g(y)[\eta_{ik} + 2hk_i k_k]dx^i dx^k, \quad (23)$$

where functions $f(y)$, $g(y)$ and h are given by (6), (5) and (18), and vector field k_i is determined by expressions (19) and (20).

In the region of parameters corresponding to spinning particles the black hole horizons disappear and a region of the rotating disk-like source is opened. The structure of this source should possess some properties of OCD-confinement and represents an old problem. The predicted exotic properties of the matter of this source [12] do not allow to construct them in four dimensions from a known sort of classical matter. The conjectured AdS/CFT correspondence gives a new approach to this problem and stimulates consideration of new models, in particular, of a bag-like models resembling the cosmic bubble models. In this case the bag-like source based on the supersymmetric domain wall models [13] can contain the AdS-region of a false vacuum inside the bag separated from the true vacuum of the outer region by a thin domain wall. As a field model, the supersymmetric version of $U(1) \times U(1)$ Witten model [14] seems the most appropriate, since it provides the long range electromagnetic field out of core. A hypothetical mechanism of the formation of the bag-like Kerr source can be connected with a phase transition governed by the value of dilaton near the core.

The AdS-BH solution of another sort was considered in [15].

Acknowledgement

I am thankful to Sergei Odintsov for paying my attention to this area and for useful discussions.

²It can be also expressed as $Y = (z - ia - \tilde{r})/(x - iy)$, where $\tilde{r} = r + ia \cos \theta$ is complex radial distance.

Appendix

The connection coefficients are (no summation over d)

$$\Gamma_{dd}^d = \frac{f'}{2f}, \quad (24)$$

$$\Gamma_{id}^d = 0, \quad (25)$$

$$\Gamma_{di}^d = 0, \quad (26)$$

$$\Gamma_{ik}^d = -\frac{g' g_{ik}}{2f}, \quad (27)$$

$$\Gamma_{dk}^i = \Gamma_{kd}^i = -\frac{g' \delta_k^i}{2g}, \quad (28)$$

$$\Gamma_{dd}^i = 0, \quad (29)$$

$$\Gamma_{jk}^i = \hat{\Gamma}_{jk}^i. \quad (30)$$

Here $\hat{\Gamma}_{jk}^i$ are connections to d-dimensional metric $\hat{g}_{ik}$.

We have also the relations

$$\Gamma_{d\mu}^\mu = \Gamma_{\mu d}^\mu = \frac{f'}{2f} + d \frac{g'}{2g}, \quad (31)$$

$$\Gamma_{i\mu}^\mu = \Gamma_{\mu i}^\mu = \hat{\Gamma}_{i\mu}^\mu = \hat{\Gamma}_{\mu i}^\mu, \quad (32)$$

$$\Gamma_{d\nu}^\mu \Gamma_{d\mu}^\nu = \left(\frac{f}{2f}\right)^2 + d \left(\frac{g'}{2g}\right)^2, \quad (33)$$

$$\Gamma_{i\nu}^\mu \Gamma_{k\mu}^\nu = \hat{\Gamma}_{ij}^l \hat{\Gamma}_{kl}^j - g_{ik} \frac{(g')^2}{2gf}, \quad (34)$$

$$\Gamma_{dd}^\mu \Gamma_{\nu\mu}^\nu = \frac{f'}{2f} \left(\frac{f'}{2f} + d \frac{g'}{2g}\right), \quad (35)$$

$$\Gamma_{ik}^\mu \Gamma_{\nu\mu}^\nu = \hat{\Gamma}_{ik}^j \hat{\Gamma}_{lj}^l - g_{ik} \frac{g'}{2f} \left(\frac{f'}{2f} + d \frac{g'}{2g}\right). \quad (36)$$

References

- [1] E. Witten, Adv. Theor. Math. Phys. **2** (1998) 253, hep-th/9802150.
 J. Maldacena, Adv. Theor. Math. Phys. **2** (1998) 231, hep-th/9711200.
 S.S. Gubser, I.R. Klebanov and A.M. Polyakov, Phys. Lett. **B428** (1998)105, hep-th/9802109;
- [2] I.R. Klebanov and A.A. Tseytlin, *Asymptotic freedom and infrared behaviour in the type 0 string approach to gauge theories*, hep-th/9812089;

- [3] J.A. Minahan, JHEP bf 9901(1999)020, hep-th/9811156; *Asymptotic freedom and confinement from type 0 string theory*, hep-th/9902074.
- [4] G. Ferretti and D. Martelli, *On the construction of gauge theories from non-critical type 0 strings*, hep-th/9811208; E. Álvarez and C. Gómez, *Non-critical confining strings and the renormalization group*, hep-th/9902012.
- [5] A.Kehagias and K. Sfetsos, Phys.Lett.**B454**(1999)270, hep-th/9902125.
- [6] A.Kehagias and K. Sfetsos, Phys.Lett.**B456**(1999)22, hep-th/9903109.
- [7] H.Liu and A.A. Tseytlin, *D3-brane – D-instanton configuration and $\mathcal{N} = 4$ super YM theory in constant self-dual background*, hep-th/9903091;
- [8] R. de Mello Koch, A. Paulin-Campbell and J. P. Rodrigues, *Non-holomorphic Corrections from Threebranes in F Theory*, hep-th/9903029;
- [9] S. Nojiri and S.D. Odintsov, *Running gauge coupling and quark-antiquark potential from dilatonic gravity*, hep-th/9904036; *Curvature dependence of running gauge coupling and confinement in AdS/CFT correspondence*, hep-th/9905200;
- [10] S. Nojiri and S.D. Odintsov Phys. Lett. **B449** (1999) 39, hep-th/9812017.
- [11] Carter B., Phys. Rev. **174**(1968) 1559
 Burinskii A., Sov.Phys.JETP **39**(1974) 193;
 Sen A. Modern Phys. Lett. **A 10**(1995)2081; Nucl.Phys **B46** (Proc.Suppl.) (1996)198;
 Burinskii A.,
 Phys.Lett. **A 185**(1994)441; Phys.Rev.**D52**(1995)5826;**D57**(1998)2392;
 Class.Quant. Grav. **16** (1999) 3497, hep-th/9903032;
 Dabholkar A., Gauntlett J., Harvey J. and Waldram D., Nucl. Phys. **B 474**(1996) 85;

- [12] Israel W., Phys. Rev. **D2** (1970) 641;
Lòpez C.A., Phys. Rev. **D30** (1984) 313;
Burinskii A. Phys.Lett.**B216**(1989)123;
- [13] Morris J.R. and Baseia D. Phys.Rev.**D 54**(1996)5217;
Cvetic M., Griffies S. and Rey S.-J. Nucl.Phys.**381**(1992)301;
- [14] Morris J.R., Phys.Rev.**D 53**(1996)2078;
- [15] S. Nojiri and S.D. Odintsov, *Running gauge coupling and quark-antiquark potential in non-SUSY gauge theory at finite temperature from IIB SG/CFT correspondence*, hep-th/9906216;